


Failed Back & Neck Surgery

What it is

Some people still have pain after back or neck surgery. Doctors sometimes call this failed back or neck surgery syndrome. It does not mean anyone did something wrong. There are still options that may help.



Common symptoms

- Back or neck pain that persists after surgery
- Leg or arm pain, burning or numbness
- Pain that returns after a period of improvement
- Reduced ability to walk, sit or work

Self-care tips

- Stay as active as you can. Short, easy walks may help.
- Change position often. Try not to sit for more than 30 to 45 minutes at a time.
- Heat or ice for 15 to 20 minutes may ease pain. Put a towel between it and your skin.
- Lift with your legs, keep the load close to your body, and do not twist.

What to expect at GSMC

At your visit: We ask about your symptoms and examine you.

Tests: You may need X-rays, an MRI or other tests. Not everyone needs them.

Your plan: We talk about your options, starting with the simplest steps. We decide together.

When to call us

- Pain that keeps getting worse, or does not get better with care
- New numbness, tingling or weakness in an arm or leg
- Pain with fever, chills or weight loss you cannot explain
- Pain that keeps you from sleeping or doing daily tasks

Call 911 or go to the emergency room for loss of bladder or bowel control, numbness in the groin or inner thighs, or sudden leg weakness.

Call 732-202-3000

One number for all offices. Questions about this handout? We are glad to help.

Our 7 offices: Whiting · Toms River (Hooper Ave) · Brick · Barnegat · Manahawkin · Freehold · West Long Branch

gsmedicalcenter.org/conditions/failed-back-neck-surgery/



Scan for the full web page

This handout is general information and is not a substitute for medical advice.